### Food and Drug Administration, HHS

tuberculosis and provides epidemiological information on this disease. *Mycobacterium tuberculosis* is the common causative organism in human tuberculosis, a chronic infectious disease characterized by formation of tubercles (small rounded nodules) and tissue necrosis (destruction), usually occurring in the lung.

(b)  ${\it Classification}$ . Class I (general controls).

## §866.3375 Mycoplasma spp. serological reagents.

(a) Identification. Mycoplasma spp. serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to Mycoplasma spp. in serum. Additionally, some of these reagents consist of Mycoplasma spp. antisera conjugated fluorescent with dve (immunofluorescent reagents) used to identify Mycoplasma spp. directly from clinical specimens. The identification aids in the diagnosis of disease caused by bacteria belonging to the genus Mycoplasma and provides epidemiological information on diseases caused by these microorganisms. Mycoplasma spp. are associated with inflammatory conditions of the urinary and respiratory tracts, the genitals, and the mouth. The effects in humans of infection with Mycoplasma pneumoniae range from inapparent infection to mild or severe upper respiratory disease, ear infection, and bronchial pneumonia.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 65 FR 2311, Jan. 14, 2000]

### §866.3380 Mumps virus serological reagents.

(a) Identification. Mumps virus serological reagents consist of antigens and antisera used in serological tests to identify antibodies to mumps virus in serum. Additionally, some of these reagents consist of antisera conjugated with a fluorescent dye (immunofluorescent reagents) used in serological tests to identify mumps viruses from tissue culture isolates derived from clinical specimens. The

identification aids in the diagnosis of mumps and provides epidemiological information on mumps. Mumps is an acute contagious disease, particularly in children, characterized by an enlargement of one or both of the parotid glands (glands situated near the ear), although other organs may also be involved.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

 $[47\ FR\ 50823,\ Nov.\ 9,\ 1982,\ as\ amended\ at\ 65\ FR\ 2311,\ Jan.\ 14,\ 2000]$ 

#### §866.3390 Neisseria spp. direct serological test reagents.

(a) Identification. Neisseria spp. direct serological test reagents are devices that consist of antigens and antisera used in serological tests to identify Neisseria spp. from cultured isolates. Additionally, some of these reagents consist of Neisseria spp. antisera conjugated with a fluorescent dye (immunofluorescent reagents) which may be used to detect the presence of Neisseria spp. directly from clinical specimens. The identification aids in the diagnosis of disease caused by bacteria belonging to the genus Neisseria. such as epidemic cerebrospinal meningitis, meningococcal disease, and gonorrhea, and also provides epidemiological information on diseases caused by these microorganisms. The device does not include products for the detection of gonorrhea in humans by indirect methods, such as detection of antibodies or of oxidase produced by gonococcal organisms.

(b) Classification. Class II (performance standards).

#### §866.3400 Parainfluenza virus serological reagents.

(a) Identification. Parainfluenza virus serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to parainfluenza virus in serum. The identification aids in the diagnosis of parainfluenza virus infections and provides epidemiological information on diseases caused by these viruses. Parainfluenza viruses cause a

#### § 866.3405

variety of respiratory illnesses ranging from the common cold to pneumonia.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 54 FR 25047, June 12, 1989; 66 FR 38792, July 25, 2001]

### §866.3405 Poliovirus serological reagents.

(a) Identification. Poliovirus serological reagents are devices that consist of antigens and antisera used in serological tests to identify antibodies to poliovirus in serum. Additionally, some of these reagents consist of poliovirus antisera conjugated with a fluorescent dye (immunofluorescent reagents) used to identify polioviruses from clinical specimens or from tissue culture isolates derived from clinical specimens. The identification aids in the diagnosis of poliomyelitis (polio) and provides epidemiological information on this disease. Poliomyelitis is an acute infectious disease which in its serious form affects the central nervous system resulting in atrophy (wasting away) of groups of muscles, ending in contraction and permanent deformity.

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 65 FR 2312, Jan. 14, 2000]

# §866.3410 Proteus spp. (Weil-Felix) serological reagents.

(a) Identification. Proteus spp. (Weil-Felix) serological reagents are devices that consist of antigens and antisera, including antisera conjugated with a fluorescent dye (immunofluorescent reagents), derived from the bacterium Proteus vulgaris used in agglutination tests (a specific type of antigen-antibody reaction) for the detection of antibodies to rickettsia (virus-like bacteria) in serum. Test results aid in the diagnosis of diseases caused by bacteria belonging to the genus Rickettsiae and provide epidemiological information on these diseases. Rickettsia are generally transmitted by arthropods (e.g., ticks

and mosquitoes) and produce infections in humans characterized by rash and fever (e.g., typhus fever, spotted fever, Q fever, and trench fever).

(b) Classification. Class I (general controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to the limitations in §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 54 FR 25047, June 12, 1989; 66 FR 38792, July 25, 2001]

#### §866.3415 Pseudomonas spp. serological reagents.

(a) Identification. Pseudomonas spp. serological reagents are devices that consist of antigens and antisera, including antisera conjugated with a fluorescent (immunofluorescent reagents), used to identify Pseudomonas spp. from clinical specimens or from cultured isolates derived from clinical specimens. The identification aids in the diagnosis of disease caused by bacteria belonging to the genus Pseudomonas. Pseudomonas aeruginosa is a major cause of hospital-acquired infections, and has been associated with urinary tract infections, eye infections, burn and wound infections, blood poisoning, abscesses, and meningitis (inflammation of brain membranes). Pseudomonas pseudomallei causes melioidosis, chronic pneumonia.

(b) Classification. Class II (special controls). The device is exempt from the premarket notification procedures in subpart E of part 807 of this chapter subject to §866.9.

[47 FR 50823, Nov. 9, 1982, as amended at 63 FR 59227, Nov. 3, 1998]

## §866.3460 Rabiesvirus immunofluorescent reagents.

(a) Identification. Rabiesvirus immunofluorescent reagents are devices that consist of rabiesvirus antisera conjugated with a fluorescent dye used to identify rabiesvirus in specimens taken from suspected rabid animals. The identification aids in the diagnosis of rabies in patients exposed by animal bites and provides epidemiological information on rabies. Rabies is an acute infectious disease of the central nervous system which, if undiagnosed, may be fatal. The disease